SPOT ANTHRACNOSE OF FLOWERING DOGWOOD

S. A. Alfieri, Jr.

Dogwood (Cornus florida L.) is one of the most popular ornamental deciduous trees in Florida. It blooms in the spring with attractive, white or pink, showy bracts. The range of habitat includes the eastern half of the United States, overlapping into portions of Mexico and Canada.

One of the more serious diseases of flowering dogwood is spot anthracnose, caused by Elsinoe corni Jenkins and Bitanc. with Sphaceloma as its imperfect stage (6). The fungus attacks stems, leaves, petioles, peduncles, bracts, and fruits (6,8). The fungus overwinters in the lesions of infected stems and initiates new infections in the ensuing new growth of flowers, leaves, and twigs (10). Rainy or humid weather is believed to be an important contributing factor to disease severity (5).

Spot anthracnose of flowering dogwood has been reported from Delaware (2,4), Florida (10,12), Georgia (9), Louisiana (3), Maryland (7), and North and South Carolina (1). During the spring of 1970 the disease was especially severe on the leaves and bracts of flowering dogwood in Gainesville, Florida.

SYMPTOMS. Spots on the leaves are circular to irregular, rather small (1-2 mm in diameter), but very noticeable. They are reddish purple with tan centers and diffuse yellow borders (Fig. 1A). The spots on older leaves have very little to no yellow borders. Spots on the bracts (Fig. 1B), petioles, peduncles, and stems are similar to the leaf spots (6,8). On the leaves and bracts the spots may be few and scattered to very abundant. Heavy infection of the leaves causes them to be reduced in size and somewhat malformed and distorted.




Fig. 1. Spot anthracnose of flowering dogwood: A) on leaves; B) on bracts.

Contribution No. 281, Bureau of Plant Pathology.

CONTROL. In 1952 Cox and Heuberger (2) in preliminary trials directed toward control of spot anthracnose of flowering dogwood have shown that captan (Orthocide 406) at $2\ lb/100\ gal$ or maneb (Manzate) at $1\ locome{1/2}\ lb/100\ gal$ provided promising results. In addition, maneb and zineb are recommended for the spot anthracnose of Viola (pansies) and copper for this same disease on avocado (11). Any of these fungicides should be applied according to the manufacturer's recommendations.

Literature Cited

- 1. Boyce, J. S., Jr. 1952. Records of flowering dogwood spot anthracnose in North and South Carolina, 1952. In Flowering dogwood spot anthracnose and its control. Plant Dis. Reptr. 36:296.
- 2. Cox, R. S. and J. W. Heuberger. 1952. Preliminary information on the use of fungicides for the control of spot anthracnose of flowering dogwood. In Flowering dogwood spot anthracnose and its control. Plant Dis. Reptr. 36: 298-299.
- 3. Driver, C. H. and A. G. Plakidas. 1953. Spot anthracnose on flowering dogwood in Louisiana. Plant Dis. Reptr. 37:448.
- 4. Jehle, R. A. and Anna E. Jenkins. 1949. Roadside survey for spot anthracnose of flowering dogwood in Maryland, Virginia, Delaware, and West Virginia. Plant Dis. Reptr. 33:198-201.
- 5. Jehle, R. A. and Anna E. Jenkins. 1952. Known distribution of flowering dogwood spot anthracnose in Maryland, May 1952. In Flowering dogwood spot anthracnose and its control. Plant Dis. Reptr. 36:296-297.
- 6. Jenkins, Anna E. 1948. Diagnosis of the Elsinoe of flowering dogwood. Wash. Acad. Sci. J. 38:362-365.
- 7. Jenkins, Anna E. 1952. Spot anthracnose of flowering dogwood found in Nassa-wanda Forest, Maryland, and isolation of the pathogen. In Flowering dogwood spot anthracnose and its control. Plant Dis. Reptr. 36:294.
- 8. Jenkins, Anna E. and A. A. Bitancourt. 1948. A spot anthracnose of flowering dogwood. Plant Dis. Reptr. 32:253-255.
- 9. Miller, J. H. 1952. Spot anthracnose (Elsinoe corni) on Cornus florida in Georgia in 1952. In Flowering dogwood spot anthracnose and its control.~ Plant Dis. Reptr. 36:295-296.
- 10. Owen, J. H. 1955. Weather as related to plant disease development in the Gainesville area of Florida from January to April, 1955. Plant Dis. Reptr. 39:467-469.
- 11. Pirone, P. P., B. O. Dodge, and H. W. Rickett. 1960. Diseases and pests of ornamental plants. Ronald Press, New York. 776 p.
- 12. Weber, G. F. 1952. Elsinoe corni Jenkins & Bitanc. on flowering dogwood in Gainesville, Florida. In Flowering dogwood spot anthracnose and its control. Plant Dis. Reptr. 36:294-295.